

Chain Inflation via Rapid Tunneling in the Landscape

Katherine Freese*

Michigan Center for Theoretical Physics, University of Michigan, Ann Arbor, MI 48109-1040

James T. Liu†

*Michigan Center for Theoretical Physics, University of Michigan, Ann Arbor, MI 48109-1040 and
Kavli Institute for Theoretical Physics, University of California, Santa Barbara, CA 93106*

Douglas Spolyar‡

Physics Department, University of California, Santa Cruz, CA 95060

Chain inflation takes place in the string theory landscape as the universe tunnels rapidly through a series of ever lower energy vacua such as may be characterized by quantized changes in four form fluxes. The string landscape may be well suited to an early period of rapid tunneling, as required by chain inflation, followed by a later period of slow tunneling, such as may be required to explain today's dark energy and small cosmological constant. Each tunneling event (which can alternatively be thought of as a nucleation of branes) provides a fraction of an e -folding of inflation, so that hundreds of tunneling events provide the requisite amount of inflation. A specific example from M-theory compactification on manifolds with non-trivial three-cycles is presented.

In 1981, Guth [1] proposed an inflationary phase of the early universe to solve the horizon, flatness, and monopole problems of the standard cosmology. During inflation, the Friedmann equation

$$H^2 = 8\pi G_4 \rho / 3 + k/a^2 \quad (1)$$

is dominated on the right hand side by a (nearly constant) false vacuum energy term $\rho \simeq \rho_{vac} \sim \text{constant}$. The scale factor of the Universe expands superluminally, $a \sim t^p$ with $p > 1$. Here $H = \dot{a}/a$ is the Hubble parameter. With sufficient inflation, roughly 60 e -folds, the cosmological shortcomings are resolved. Inflationary models fall into two classes: tunneling models such as Guth's original model and rolling models where a scalar field slowly rolls down an extremely flat potential.

In Chain Inflation [2], the universe tunnels rapidly through a series of vacua separated from one another by energy barriers. One can imagine a multidimensional potential containing many "bowls" or minima of varying energies. The universe starts out in a high-energy minimum, and then sequentially tunnels down to bowls of ever lower energy until it reaches the bottom. During the time spent in any one of these bowls, the universe inflates by a fraction of an e -fold. After many hundreds or thousands of tunneling events, the universe has inflated by the requisite 60 (or so) e -folds to resolve the cosmological problems outlined above, as well as satisfying $\delta\rho/\rho$ [3]. At each stage, the phase transition is rapid enough that percolation is complete, though thermalization and reheating does not take place until the final few stages of tunneling near the bottom of the potential. In principle the energy scales of the potential can vary widely, from GeV scale up to almost Planck scale. No fine-tuning of the shape of the potential is required (the potential need not be flat).

A simple way to visualize Chain Inflation is to imagine

a tilted cosine potential (sometimes known as a "washboard potential") as the effective path through the multidimensional landscape. Then the universe tunnels from a high energy minimum of the cosine down to the next minimum and so on. In fact, we showed in [4] that the QCD axion, which has been proposed as a byproduct of a solution to the strong CP problem in the theory of strong interactions, can have a tilted cosine potential and hence is a candidate for the inflaton in chain inflation; in this case inflation takes place at the QCD scale and can in principle be experimentally tested in the foreseeable future.

In this paper we focus instead on chain inflation in the string landscape. Chain inflation is quite generic in the string landscape, as the universe tunnels through a series of metastable vacua, each with different fluxes[23]. There has been a great deal of interest in the fact that the vacuum of string theory is not unique; in fact there appear to be at least 10^{200} different vacua. Even with compactification on a specific Calabi-Yau manifold, there is still a large variety of background fluxes that could be turned on, providing a huge number of different possible vacua. These vacua with different fluxes are disconnected in the multidimensional landscape, with barriers in between them. Thus the only way to proceed from one to the other is via tunneling. Chain inflation is the result of rapid tunneling through a series of these vacua.

Chain inflation. In chain inflation, the universe begins in some metastable minimum of the potential with a positive value of vacuum energy. From this minimum there would presumably be a variety of possible directions in field space in which the system could tunnel to a lower minimum. However, many of the neighboring minima would be inaccessible because the tunneling rate into them is too slow; instead, the system would tunnel more quickly in a different direction. The system thus chooses

the “path of least resistance”, *i.e.* it moves through a series of minima to which the tunneling is the most rapid.

From a given initial minimum, tunneling will proceed to the neighboring minimum with the largest drop in energy, lowest barrier height and smallest difference in field value. In our first paper, (hereafter Paper 1) [2], we quantified this statement by modeling any single tunneling event using a scalar field with an asymmetric double well potential

$$V(\phi) = \frac{1}{4}\lambda(\phi^2 - a^2)^2 + \frac{\epsilon}{2a}(\phi - a). \quad (2)$$

Then the tunneling rate is determined by three quantities: the barrier height $\lambda a^4/4$, energy difference between vacua ϵ , and width of the potential $2a$. We noted that, for the case where all three scales are comparable, the tunneling rate is on the borderline of being extremely rapid or extremely slow relative to a Hubble time. The reason for this extreme sensitivity to parameters is the exponential dependence of the tunneling rate on the bounce action, $\Gamma \sim e^{-B}$ where (in the thin-wall approximation) [5, 6, 7]

$$B = \frac{27\pi^2}{2} \frac{\tau^4}{|\epsilon|^3}. \quad (3)$$

Here ϵ is the drop in energy and τ is the tension of the brane interpolating between the two vacua. In particular, for the above $\lambda\phi^4$ potential, the brane tension is given by $\tau = \frac{2\sqrt{2}}{3}\sqrt{\lambda}a^3$. Hence, for natural parameter choices, the tunneling can be either very fast or very slow. The first choice of the field will be to move in the fastest direction of large ϵ , small barrier height, and small width a .

The string landscape. A typical picture of the string landscape involves a complicated distribution of numerous metastable minima with very small energy differences and separated by large (*i.e.* string scale) barrier heights. This view of the landscape has been driven in part by attempts to solve the cosmological constant problem, in which case one is naturally driven to consider energy differences on the order of the presently observed vacuum energy, $\epsilon \sim (10^{-3} \text{ eV})^4$. From this point of view, it is generally expected that tunneling would be extremely slow. However, we note that, although metastable minima with very small energy differences are naturally present in the landscape, they would generically be well separated from each other. Instead, adjacent minima are much more likely to have large energy differences for the simple reason that the natural scale for a single configuration change (*e.g.* the nucleation of a D-brane instanton) is given by the string scale, and *not* by something like 10^{-3} eV . Since both energy differences and barriers between adjacent minima in the landscape are expected to be characterized by the same string scale, this indicates that the tunneling rate may be quite large. Furthermore, Tye recently noted another mechanism for

fast tunneling [8]: even in situations where the tunneling between neighboring minima is typically slow, resonant effects may greatly enhance the tunneling rates.

Portions of the landscape can be quite well suited to an early period of rapid tunneling, as required by chain inflation, followed by a later period of slow tunneling, such as may be required to explain today’s dark energy and small cosmological constant. As we noted in Paper 1, given a choice of tunneling to a number of lower energy vacua of different energies, the field will take the largest steps first. The field will tunnel in the direction of the larger vacuum energy difference, as the tunneling in that direction is fastest. For example, if the field has a choice of tunneling into a minimum with energy difference $\epsilon \sim (0.1 m_{pl})^4$ or a minimum with energy difference $\epsilon \sim (10^{-10} m_{pl})^4$, it will choose the former. The path taken by the field quite naturally lends itself to an early period of chain inflation with rapid tunneling. Subsequently, the field slows down, as the only remaining minima it can tunnel to have smaller and smaller energy differences from its current position.

We also note that chain inflation is *not* a model of eternal inflation. During the rapid tunneling phase, the rate of tunneling back up the potential is very small. Chain inflation is a dynamical model in which the field chooses the downward path of most rapid tunneling. Hence, the universe is driven universally to one or at most a few final states, rather than a huge number of them. Given any minimum the field is sitting in, there may be more than one rapid path, but not a huge number of them. This arbitrariness of path must be considered when evaluating density perturbations in the model. One may need to invoke the anthropic principle to explain our current universe, but only to a small degree.

Tunneling and the cosmological constant. While chain inflation was initially formulated in terms of scalar fields [2], it may equally well occur in any landscape which allows rapid tunneling through a chain of metastable minima. Since typical string models populate the landscape by turning on various fluxes, it is natural in this case to have flux-driven chain inflation (and in particular four form flux when viewed in 3+1 dimensions). Here we will see that the energy difference between minima indeed decreases as one goes to lower energy vacua, so that we can have the tunneling rate change from initially being rapid to subsequently becoming smaller. It is possible that this change in tunneling rate leads from a period of chain inflation to a universe stuck in its current small value of the cosmological constant.

To compute the tunneling, we follow the lead of Brown and Teitelboim [9, 10] (hereafter BT). The BT mechanism may be motivated by the observation that, in 1+1 dimensional electrodynamics, an electron/positron pair can be produced in the presence of a sufficiently strong background electric field. The particles are physically pulled apart by the original field, and the region between

them has a reduced value of electric field. The world lines of the electron and the positron can be thought of as membranes (domain walls), with the electric fluxes in the regions interior to the membranes reduced by a quantized amount. One can think of this process as the nucleation of membranes in the presence of background electric fields. A series of nucleation events takes place, each time reducing the value of the electric flux by a quantized amount.

As in BT, we are interested in the four-dimensional generalization of this process, in which background fields cause bubble nucleation of lower energy regions with lower field values. While the BT mechanism relies on four-form fluxes in $3+1$ dimensions, such fluxes have numerous potential origins in the string theory landscape. For example, in the case of IIB flux vacua, both NSNS and RR three-form internal flux are generically present. Taking the Hodge dual results in seven-form flux, which reduces on three-cycles of the internal Calabi-Yau manifold to yield four-form fluxes in $3+1$ dimensions. This flux may then be reduced through the nucleation of both wrapped NS5-branes and wrapped D5-branes. Alternatively, from an M-theory (or IIA) point of view, the M-theory four-form can take values in either space-time or internal space, with corresponding flux reduced through the nucleation of either M2-branes or wrapped M5-branes, respectively. In all such cases, the nucleated branes manifest themselves as membranes in the $3+1$ dimensional spacetime.

The nucleation of domain walls changes the value of the vacuum energy by giving rise to quantized drops in the value of the background fluxes; four-form field strengths play the role of a dynamical vacuum energy in four dimensional spacetime,

$$\Lambda = \Lambda_{\text{bare}} + \frac{1}{2}F_4^2, \quad (4)$$

where Λ_{bare} is the bare cosmological constant, and $F_{\mu\nu\rho\sigma} = F_4\epsilon_{\mu\nu\rho\sigma}$, so F_4 is the magnitude of a four-form field which can change its value through the nucleation of membranes. The change of the field strength across the membrane is

$$\Delta F_4 = \pm q, \quad (5)$$

where the charge q is constant so that the flux values are quantized, at least in string theory. Hence, Eq. (4) becomes

$$\Lambda = \Lambda_{\text{bare}} + \frac{1}{2}n^2q^2, \quad (6)$$

where the integer n labels the integer flux.

At each stage of domain wall production, the integer flux changes from $n \rightarrow n-1$, and the vacuum energy changes by an amount

$$\epsilon = -(n-1/2)q^2. \quad (7)$$

This is the energy difference between vacua in chain inflation; one can think of this process as tunneling from one vacuum to another whose energy is lower by the amount given in Eq. (7).

Bousso and Polchinski (hereafter BP) extended the BT approach to include multiple four-form fluxes [11]. They noted that the quantized energy changes in Eq. (7) with a single flux in BT are too large to end up with a small vacuum energy today of order the dark energy, $\Lambda \sim 10^{-120}m_{pl}^4$. On the other hand, compactification manifolds could easily have a large number of cycles, and turning on fluxes on all these cycles leads to many effective four-form fluxes in $3+1$ dimensions. By using a large number J of fluxes with a J -dimensional grid of vacua, they noted that the spectrum of allowed values of Λ can then be sufficiently dense to explain the observed value of the cosmological constant in some portion of the universe. We will follow their multi-flux proposal, but dispense with subscripts that differentiate between one flux and another (e.g. we will use the notation n to generically mean the integer flux for any one of the four forms). Furthermore, while BP had the goal of neutralizing the cosmological constant in this fashion, we are more directly interested in chain inflation in the landscape. Hence we do not necessarily concern ourselves with the details of the endpoint of the BP scenario.

Requirements for chain inflation. In the zero temperature limit, the nucleation rate Γ per unit spacetime volume for producing bubbles of true vacuum in the sea of false vacuum through quantum tunneling can be expressed as

$$\Gamma(t) = Ae^{-S_E} = Ae^{-(B-S_{BG})}. \quad (8)$$

Here the Euclidean action S_E is the difference between the instanton action B given (in the thin-wall limit) by Eq. (3) and the background Euclidean action of de Sitter space $S_{BG} = -8\pi^2/H^2$. A is a determinantal factor which can be estimated as $A \sim 1$ [12, 13]. As seen in Eq. (3), the instanton action depends on the difference in energy ϵ between the inside and outside the bubble and on the domain wall tension τ . Chain inflation consists of a number of rapid tunneling steps, in which case the bubbles are small compared to the horizon size. In this case, the gravitational correction S_{BG} can be ignored, so that $S_E \sim B$. Later we will comment on a possible treatment beyond the thin-wall limit.

Guth and Weinberg [14] have shown that the probability of a point remaining in a false de Sitter vacuum is approximately

$$p(t) \sim \exp\left(-\frac{4\pi}{3}\beta Ht\right), \quad (9)$$

where the dimensionless quantity β is defined by $\beta \equiv \Gamma/H^4$. The number of e -foldings per tunneling event is

$$\chi = \int Hdt \sim H\tau = \frac{3}{4\pi\beta}. \quad (10)$$

The authors of [14, 15] calculated that a critical value of

$$\beta \geq \beta_{\text{crit}} = 9/4\pi \quad (11)$$

is required in order to achieve percolation and thermalization. In terms of number of e -foldings, this is

$$\chi \leq \chi_{\text{crit}} = 1/3. \quad (12)$$

As long as this is satisfied, the phase transition at each stage takes place quickly enough so that ‘graceful exit’ is achieved. Bubbles of true vacuum nucleate throughout the universe at once, and are able to percolate. Sufficient inflation requires the total number of e -foldings to satisfy

$$N_{\text{tot}} > 60. \quad (13)$$

Old inflation [1] failed in that a single tunneling event can satisfy either Eq. (12) or Eq. (13) but not both. Chain inflation satisfies both equations by having a chain of tunneling events, each of which provides only a fraction of an e -folding but adding up to a total of at least 60-folds.

Chain inflation in string theory. Given the above requirements, we now examine the feasibility of realizing chain inflation in the string landscape picture. In general, fluxes in string theory are closely related to branes, in that p -branes are natural sources for $(p+2)$ -form fluxes. The model we have in mind is essentially that of BP [11], where we start with a D dimensional theory (where D may be 10 or 11) compactified down to $3+1$ dimensional spacetime. The objects that we are interested in are BPS p -branes which satisfy a D dimensional relation between charge and tension

$$\mu_p^2 = 2\kappa_D^2 \tau_p^2. \quad (14)$$

For Dp -branes we have $\tau_p = (\sqrt{\pi}/\kappa_{10})(2\pi\sqrt{\alpha'})^{3-p}$, while for M2 and M5-branes, we have instead $\tau_2 = 2\pi M_{11}^3$ and $\tau_5 = 2\pi M_{11}^6$, respectively, where $2\pi M_{11}^9 = 1/(2\kappa_{11}^2)$.

In order to obtain four-form fluxes in $3+1$ dimensions, we consider p -branes wrapped on $(p-2)$ -cycles of the compactification manifold. In this case, the effective brane tension in four dimensions is given simply by

$$\tau = \tau_p V_{p-2}, \quad (15)$$

where V_{p-2} is the volume of the corresponding $(p-2)$ -cycle which the brane wraps. In addition, the brane charge measured in four dimensions can be shown to be

$$q = \frac{\mu_p V_{p-2}}{\sqrt{V_{D-4}}}, \quad (16)$$

where V_{D-4} is the volume of the compactification manifold. Using Eqs. (15) and (16), we see that the D -dimensional BPS relation, Eq. (14), reduces to the universal four-dimensional relation

$$\tau = \frac{1}{\sqrt{2}} m_{\text{pl}} q, \quad (17)$$

where the four-dimensional Planck mass ($m_{\text{pl}} = 2.43 \times 10^{18}$ GeV) is related to the D -dimensional one by

$$m_{\text{pl}}^2 \equiv \frac{1}{8\pi G_4} = \frac{V_{D-4}}{\kappa_D^2}. \quad (18)$$

Although we view this model of $(p+2)$ -form fluxes reduced to four form fluxes on various cycles as a toy model of the string landscape, we believe that it captures many of the generic features common to any realistic flux compactification setup. In particular, we expect that the effective tensions and charges are given, at least within the correct order of magnitude, by Eq. (17).

Starting for a given metastable minimum specified by a particular set of fluxes, we now estimate the tunneling rate of Eq. (8), or equivalently, the membrane nucleation rate. Although a generic state in the landscape involves many independent fluxes (specified by a set of integers $\{n_i\}$), because they are independent (at least to leading order), it is sufficient to consider the nucleation of a single membrane, which changes the flux of one of the four-forms from n to $n-1$. In this case, the change in the vacuum energy is given by Eq. (7). Taking $n \gg 1$, we have

$$|\epsilon| \approx nq^2. \quad (19)$$

For $n \gg 1$, and for Λ sufficiently large[24], gravity can be neglected so the bounce action is essentially given by Eq. (3). Inserting τ and ϵ corresponding to the nucleated membrane into this expression yields [9, 11]

$$B_n = \frac{27\pi^2 m_{\text{pl}}^6}{16n^3 \tau^2}, \quad (20)$$

where we have used the BPS relation, Eq. (17), to relate the brane charge q with its tension τ . We note that the semiclassical approximation in which the instanton calculations have been computed is valid for $B_n \gg 1$. Thus the above expression cannot be trusted in the fast tunneling regime that we are interested in (where $B_n \sim 1$). Nevertheless, the qualitative behavior of B_n as a function of the brane tension ought to remain valid, and that is mainly what we are interested in.

It ought to be apparent from Eq. (20) that tunneling between adjacent minima is governed by both the amount of flux n along a given cycle as well as the tension τ/m_{pl}^3 of the corresponding brane (in Planck units) that must be nucleated to reduce this flux. Note, in particular, that tunneling is enhanced for larger brane tensions. The reason behind this somewhat counterintuitive result is that the change in energy ϵ increases as $\epsilon \sim \tau^2$, and this is more than sufficient to compensate for the increased brane tension. As far as this toy model is concerned, these adjacent minima are well separated in energy, and hence fast tunneling is very much generic in this landscape. Metastable minima with tiny differences in energy (as envisioned in BP) are well separated, and are

only connected through multiple flux jumps along many simultaneous cycles. Tunneling between such states are highly suppressed. However, this is irrelevant, as the system naturally tunnels through the most favorable path at any given stage in the chain. In the simple case where all brane tensions are identical, the cycle with the largest n (*i.e.* supporting the most flux) is the one which preferentially undergoes tunneling. In this case, at the beginning of chain inflation, the system is in a state with fluxes specified by an initial set of independent quanta $\{n_i^{\text{initial}}\}$, but can naturally end up on the BP circle with $\{n_i^{\text{final}} \approx n_0\}$.

Of course, just because the system undergoes a series of tunneling events until it dynamically neutralizes Λ_{bare} does not in itself ensure that chain inflation actually occurs in the landscape. To see whether chain inflation is possible, we first compute the quantity $\beta = \Gamma/H^4$ using Eq. (8) for the tunneling rate as well as the Hubble relation $H^2 = \Lambda/3m_{\text{pl}}^2$. We find

$$\beta_n \equiv \frac{\Gamma}{H^4} = 9 \left(\frac{\Lambda_n}{m_{\text{pl}}^4} \right)^{-2} e^{-B_n}, \quad (21)$$

where Λ_n is given by Eq. (6) generalized to multiple fluxes

$$\Lambda_n = \Lambda_{\text{bare}} + m_{\text{pl}}^{-2} \sum_i n_i^2 \tau_i^2, \quad (22)$$

and we have introduced the subscript n to indicate that the quantities are dependent on the amount of flux that is threading the cycle that undergoes this stage of tunneling.

At any step in the chain, β_n must satisfy the inequality

$$\beta_n \geq \frac{9}{4\pi} \approx 0.716, \quad (23)$$

given by Eq. (11) in order for percolation to succeed at each tunneling transition. From Eq. (21), this puts a requirement on the bounce action

$$B_n \lesssim 2.53 - 2 \log(\Lambda_n/m_{\text{pl}}^4). \quad (24)$$

Note that Λ_n is the vacuum energy of the metastable minimum specified by n (actually the complete set $\{n_i\}$). Depending on the parameters of chain inflation, this can be anywhere from near (but below) the Planck scale down to few tens of MeV's (in order to generate a sufficient reheating temperature) [2]. Furthermore, Λ_n changes throughout the chain of tunneling events, starting from some initial value Λ_{initial} , and dropping down to $\Lambda_{\text{final}} \approx 0$ at the end of inflation.

Our final requirement is given by Eq. (13), which ensures that sufficient inflation occurs. In particular,

$$N_{\text{tot}} = \sum \frac{3}{4\pi\beta_n} \geq 60, \quad (25)$$

where the sum is over the total number of tunneling events during inflation. As chain inflation proceeds, the value of β_n given by Eq. (21) changes because of two competing effects. The first is that, as Λ_n is reduced, β_n naturally tends to increase. However, the second is that, as the overall flux is driven down, the values of n decrease. This leads to a larger bounce action B_n , which in turn leads to an exponential suppression of β_n . Because β_n is variable, Eq. (25) does not lead itself to a simple expression for the total number of tunneling events. However, a necessary condition may be given as

$$\# \text{ of tunnelings} \geq 80\pi\beta_{\text{min}} \geq 180, \quad (26)$$

where the second inequality is simply the percolation requirement of Eq. (23).

The constraints, Eqs. (24) and (26), on the bounce action and the number of tunnelings must be simultaneously satisfied in chain inflation, using the effective four-dimensional brane tension given by Eq. (15). For example, taking the M-theory example of BP, we may consider the nucleation of both M2-branes and M5-branes wrapped on three-cycles of the seven-dimensional compactification manifold. Inserting the appropriate brane tensions into Eq. (20), we obtain

$$B_n = \frac{27}{64n^3} \left(\frac{m_{\text{pl}}}{M_{11}} \right)^6 \times \begin{cases} 1 & \text{M2,} \\ (V_3 M_{11}^3)^{-2} & \text{M5.} \end{cases} \quad (27)$$

In order to arrive at a large number of four-form fluxes, we consider the case of wrapped M5-branes. In this case, one example of a satisfactory set of parameters is to take $M_{11} = 10^{-3} m_{\text{pl}}$ along with $V_3 M_{11}^3 = 10^3$. Starting at $\Lambda_{\text{initial}} = (10^{-1} m_{\text{pl}})^4$, we see that $\beta_n > 1$ for $B_n < 21$, which is obtained for $n > 2735$. Assuming inflation ends at $\Lambda_{\text{final}} = (10^{-20} m_{\text{pl}})^4$, a similar computation indicates that $n > 1045$ at the end of inflation. Note that, for these parameters, the effective wrapped brane tension is $\tau = 2\pi \times 10^{-6} m_{\text{pl}}^3$. Eq. (22) then indicates that each flux contributes on the order of $4n^2 \times 10^{-11} m_{\text{pl}}^4$ to Λ . For $n \sim 10^3$, the energy difference per step is then on the order of $\epsilon \sim 10^{-7} m_{\text{pl}}^4$. As a result, about 1000 tunneling events will be needed to neutralize Λ_{initial} .

The above requirements on n (along with that of a sufficient number of tunnelings) may be satisfied by an appropriate choice of Λ_{bare} in the BP framework. Note that such large values of flux may destabilize the back-ground manifold due to backreaction, or could prove difficult to realize because of tadpole constraints. However, this issue is unresolved (see, *e.g.*, [16, 17]). In this case, it might be better to consider the alternative of large effective brane tension τ rather than large n as responsible for rapid tunneling. In this M5-brane example, this corresponds to taking a compactification manifold with large three-cycles.

End of inflation. We have used four forms to illustrate a possible implementation in M-theory of the basic ideas

of chain inflation. The nice feature exists that, as time goes on, the tunneling rate automatically slows down as the integer units of flux n decreases. However, it is hard to see how inflation ends. While the general idea of rapid tunneling followed by slow tunneling is attractive, the trouble is that there can be an intermediate regime of medium tunneling. This intermediate regime would mimic the reheating problems of old inflation, in that the universe could end up with too many empty big bubbles that have not percolated. However, this could potentially be avoided through resonant tunneling [8] or devaluation [18]. Another possible way to avoid this problem is to have a very large Planck-scale negative value of Λ_{bare} that gets cancelled off by the positive vacuum contributions we have been discussing in this paper. Then the landscape would involve a steep set of metastable minima (analogous to a linearly tilted cosine potential) that persists into negative values of the vacuum energy. The naive thin-wall computation of tunneling would give us rapid tunneling into AdS, which would obviously be disastrous. However, it has been argued by Banks and others [19] that tunneling into AdS is forbidden, in which case chain inflation would simply stop at the last positive value of vacuum energy before going negative.

Beyond thin-wall. Our treatment of membrane nucleation as an effective tunneling process for chain inflation has up to this point assumed the thin-wall limit, where the energy difference between barriers is much smaller than the barrier height. The computation of the instanton action in Eq. (3) by [5, 6, 7] made this assumption. Banks, however, noted that one can go beyond the thin-wall computation [19]. As mentioned previously, the tunneling process is the higher dimensional analog of pair production in 1+1 dimensional electrodynamics, the massive Schwinger model. Using techniques of bosonization, this model can be shown [20] to be equivalent to a scalar field theory with Lagrangian

$$L = \frac{1}{2}[(\Delta\phi)^2 - \frac{e^2}{\pi}\phi^2 + M^2\cos(\phi)]. \quad (28)$$

In this formulation, pair production is described as vacuum decay of the false vacua at minima of the cosine, which correspond to backgrounds with fixed quantized values of electric flux.

Banks notes that in higher dimensions, though the bosonization does not generalize, one can still introduce a scalar field through p -forms:

$$F_{\mu_1\ldots\mu_p} = \epsilon_{\mu_1\ldots\mu_p}\phi. \quad (29)$$

He argues that a derivative expansion for the effective action for ϕ , with inclusion of effects of membranes, will again lead to the form of Eq. (28), though with the cosine replaced by a periodic function. In [4], we studied chain inflation with a tilted cosine potential similar to the one in Eq. (28). A treatment of tunneling beyond the thin-wall limit with a tilted cosine potential is underway [21].

Discussion. We have shown that chain inflation can take place in the string theory landscape as the universe tunnels through a series of ever lower energy vacua. This is not a variant of eternal inflation, as tunneling back up the potential is suppressed [13]. We illustrated an example in which these vacua are characterized by quantized changes in four-form fluxes. Although we have focused on effective 3+1 dimensional dynamics, these ideas could of course be generalized to a series of tunneling events via p -forms in p -dimensions.

We also note that chain inflation can take place at low scales (*e.g.* TeV) and so can be compatible with low scale gravity $M_{11} \sim \text{TeV}$ as is the case for large extra dimensions. Most single field rolling models fail at such low scales, as they require potentials with Planck scale width and GUT scale height.

A recent paper by de Alwis [22] claims that, using the string effective supergravity action together with branes, there is no tunneling. However, this result was not derived in the full string theory. Here we have assumed that the creation of M2-branes or M5-branes, which is effectively a tunneling process, makes physical sense in an estimate of the more precise theory.

Acknowledgements

We would like to thank B. Acharya, T. Banks, K. Bobkov, R. Bousso, O. DeWolfe, M. Dine, G. Kane, D. Marolf, L. McAllister, J. Polchinski and S. Watson for many useful discussions. KF thanks N. Arkani-Hamed for pointing out that chain inflation with four forms is generic in the string landscape. This work was supported in part by the US Department of Energy under grant DE-FG02-95ER40899, the National Science Foundation under grant PHY99-07949, and the Michigan Center for Theoretical Physics under grant MCTP-06-27. KF would like to thank the Miller Institute at the University of California, Berkeley, for support and hospitality, and JTL wishes to acknowledge the hospitality of the KITP.

* Electronic address: ktfreese@umich.edu

† Electronic address: jimliu@umich.edu

‡ Electronic address: dspolyar@physics.ucsc.edu

- [1] A. H. Guth, *Phys. Rev. D* **23**, 347 (1981).
- [2] K. Freese and D. Spolyar, *JCAP* **0507**, 007 (2005).
- [3] B. Feldstein and B. Tweedie, hep-ph/0611286.
- [4] K. Freese, J. Liu, and D. Spolyar, hep-ph/0502177, *Phys. Rev. D* **72**, 123521 (2005).
- [5] C. Callan and S. Coleman, *Phys. Rev. D* **16**, 1762 (1977).
- [6] S. Coleman, *Phys. Rev. D* **15**, 2929 (1977); M. B. Voloshin, I. Yu. Kobzarev, and L. B. Okun, *Yad. Fiz* **20**, 1229 (1974) [*Sov. J. Nucl. Phys.*, **20**, 644 (1975)].
- [7] S. Coleman and F. de Luccia, *Phys. Rev. D* **21**, 3305 (1980).

- [8] S.H. Tye, hep-th/0611148
- [9] J.D. Brown and C. Teitelboim, *Phys. Lett. B* **195**, 177 (1987).
- [10] J.D. Brown and C. Teitelboim, *Nucl. Phys. B* **297** 787 (1988).
- [11] R. Bousso and J. Polchinski, *JHEP* **0006**, 6 (2000).
- [12] J. Garriga, *Phys. Rev. D* **49**, 6327 (1994).
- [13] D. Schwartz-Perlov and A. Vilenkin, hep-th/0601162
- [14] A. H. Guth and E. Weinberg, *Nucl. Phys. B* **212**, 321 (1983).
- [15] M.S. Turner, E.J. Weinberg, and L.M. Widrow, *Phys. Rev. D* **46**, 2384 (1992).
- [16] O. DeWolfe, A. Giriyavets, S. Kachru and W. Taylor, *JHEP* **0507**, 066 (2005) [hep-th/0505160].
- [17] J. Kumar and J. D. Wells, *JHEP* **0509**, 067 (2005) [hep-th/0506252].
- [18] K. Freese, J. T. Liu and D. Spolyar, *Phys. Lett. B* **634**, 119 (2006) [hep-ph/0510065].
- [19] T. Banks, hep-th/0211160.
- [20] S. Coleman, *Phys. Rev. D* **11**, 2088 (1975).
- [21] K. Freese, J. Liu, and D. Spolyar, in progress.
- [22] S.P. deAlwis, hep-th/0605184.
- [23] KF thanks N. Arkani-Hamed for drawing attention to this fact.
- [24] Since chain inflation occurs prior to neutralization of the cosmological constant, there is no contradiction between this assertion and the framework of BP.